## **RESTORATION: POSTERIOR AMALGAM**

25

## **2025 DENTAL THERAPY EXAM**

Toot	h #:	
MO	DO	MOD

CRITICAL ERRORS			
The restoration is fractured	No	Yes	

ACC = Adheres to Criteria SUB= Marginally Substandard **DEF=** Critically Deficient MARGIN INTEGRITY AND SURFACE FINISH Margin Excess/Deficiency A. No marginal deficiency. There is no evidence of pits and/or voids at the cavosurface margin. ACC B. Marginal excess  $\leq 0.5$  mm at the restoration-tooth interface, detectable either visually or with the tine of an explorer. A. DEFICIENCY: The restoration-tooth interface is detectable visually or with the tine of an explorer, and there is evidence of marginal deficiency  $\leq 0.5$  mm, which can include pits and/or voids at the cavosurface margin. SUB B. **EXCESS:** Any marginal excess is detectable visually or with the tine of an explorer, and the discrepancy is > 0.5 mm but ≤ 1.0 mm. A. There is evidence of marginal deficiency of > 0.5 mm which includes pits and voids at the cavosurface margin, and/or there DEF is an open margin. B. There is a marginal excess of > 1.0 mm. Adjacent Tooth Structure There is no or minimal evidence of unwarranted or unnecessary removal, modification, or recontouring of tooth structure ACC adjacent to the restoration. (Enameloplasty) DEF There is gross enameloplasty. CONTOUR, CONTACT, AND OCCLUSION Interproximal Contact Interproximal contact is visually closed, and the contact is adequate in size, shape, or position but demonstrates little resistance ACC to dental floss. A. The interproximal contact is visually open or concave/irregular, allowing for food impaction. DEF B. The interproximal contact will not allow floss to pass. **Centric/Excursive Contacts** When checked with articulating paper, all centric and excursive contacts on the restoration are consistent in size, shape, and ACC intensity with such contacts on other teeth in that quadrant. When checked with articulating paper, the restoration is in hyper-occlusion, making it inconsistent in size, shape, and intensity SUB with the occlusal contacts on surrounding teeth, and it requires adjustment. DEF There is gross hyper-occlusion so that the restoration is the only point of occlusion in that quadrant.

TREATMENT MANAGEMENT		
Adjacent Tooth Damage		
ACC	Any damage to adjacent tooth/teeth can be removed with polishing without adversely altering the shape of the contour and/or contact.	
DEF	There is evidence of gross damage and/or alteration to adjacent and/or opposing hard tissue inconsistent with the procedure.	
Soft Tissue Damage		
ACC	The soft tissue is free from damage or there is tissue damage that is consistent with the procedure.	
DEF	There is gross iatrogenic damage to the soft tissue inconsistent with the procedure and pre-existing condition of the soft tissue.	