Local Anesthesia Examinations

- Computer-based
- Patient-based Clinical

2025 CANDIDATE MANUAL



ALL RIGHTS RESERVED

American Board of Dental Examiners

ADEX Testing Media

Copyright © 2024-2025

TABLE OF CONTENTS

LOCAL ANESTHESIA EXAMINATION	1
General Information	1
COMPUTER-BASED EXAM	2
Computer-based Exam References	2
PATIENT-BASED CLINICAL EXAM	4
Equipment and Materials	4
Patient Criteria	5
Grading Criteria	7
Clinical Examination Day	11
Onsite Question and Answer Session/Clinic Tour	11
Clinic Entrance and Exam Procedures	11
Exam Completion	12
Preliminary Results Reporting	12
Optional Onsite Retake Exam	13
Patient-Based Clinical Examination Overview	15

ADDENDUM

Additions or modifications after initial publication.

 CDCA-WREB-CITA is now the American Board of Dental Examiners. CDCA-WREB-CITA, the nation's leading dental exam administrator, and the American Board of Dental Examiners (ADEX), developer of the national ADEX licensure exams, are now combined under a single entity: the American Board of Dental Examiners. This is also reflected in a new organizational logo and updated branding. (09/16/2025)

LOCAL ANESTHESIA EXAMINATION

General Information

The American Board of Dental Examiners **Local Anesthesia Examination** offers separate exam components as a pathway to licensure or certification in the administration of local anesthesia where applicable. The onus is on the Candidate to ensure that all requirements in a given state are met. Contact the state board from which local anesthesia licensure or certification is sought for their testing requirements.

- Computer-based in all jurisdictions that require completion of a written exam.
- Patient-based Clinical in only those jurisdictions that also require completion of a
 patient-based clinical exam. Note that successful completion (passing scores) of both
 Computer-based and Patient-based Clinical exam components is required to achieve an
 overall passing score. All components of the Local Anesthesia Examination series must
 be successfully completed (passing scores) within 18 months of the initial attempt of any
 component of the series.

Official Resources. Candidates are responsible for reading and understanding the Candidate Manual in conjunction with the Candidate Guide and noted materials published by the American Board of Dental Examiners. Candidates bear all risk of any misunderstanding resulting from the use of or reliance on unofficial information or materials.

- Candidate Guide. Refer to the Candidate Guide for information regarding education requirements and exam eligibility, identification requirements, candidate conduct, misconduct policies, computer-based testing accommodations, and more available at <u>ADEXtesting.org</u>.
- Candidate Manual, sections:
 - Computer-based exam.
 - Patient-based Clinical exam. Patient criteria, performance expectations, and examination forms.
- **Test Prep.** Local Anesthesia Clinical examination review video is accessible via the <u>Test</u> Prep page or the ADEX Testing Media YouTube page.
- Site Info Sheet. Site-specific information pertaining to the clinical exam site host, such
 as school-provided supplies, rental equipment, infection control protocols, etc. can be
 found in the Site Info Sheet for the respective exam site host on the ADEXtesting.org
 website.
- Submit any exam-related questions via Contact Us prior to the scheduled exam.

COMPUTER-BASED EXAM

The purpose of the computer-based Local Anesthesia Computer-based Examination is to evaluate candidate knowledge, application, and problem-solving skills regarding the safe and appropriate use of local anesthesia.

The timed examination is 60 minutes in length. Select-choice items are both discipline and case-based and have only one correct response. Up to five (5) field test questions are included, but not calculated in your final score.

Multiple test forms are used to ensure the integrity and security of the examination. To address any possible variation in difficulty level among the various examination forms, equating procedures are used to ensure candidates of comparable competency will be equally likely to pass the examination.

A maximum of 100 points is possible and final score of 75 or higher is required to pass the Local Anesthesia Computer-based examination.

Subject matter includes:

- Anatomy (head and neck) and physiology.
- Pharmacology of anesthetic agents and vasoconstrictors, clinical actions and maximum recommended doses of specific agents.
- Method of delivery of local anesthesia, including armamentarium, selection of injection type, and administration technique.
- Medical history interpretation; prevention, recognition, and management of possible complications, and life support.

Exam Appointment Scheduling. After registration processing and approval, it may take up to 24 hours or more for the green "Schedule My Computer-based Exam Appointment" ribbon to display in your online Candidates.ADEXtesting.org candidate profile. All the information necessary to schedule your computer-based examination appointment at a **Prometric Test Center** location is contained within the ProScheduler application of your online profile.

Valid Identification. Only valid, non-expired forms of signature-bearing U.S. **government-issued photo identification** (e.g. driver license, passport, military ID) are accepted at Prometric Test Center locations.

Candidates who do not have the required valid form(s) of identification risk forfeiture of their examination attempt and registration fees (see <u>Candidate Guide</u>, <u>Candidate Attendance & "No Show" Policy</u>).

Calculators. Upon check-in at the Prometric Test Center location, the candidate may request an erasable **note board**, **marker and hand-held calculator**, if necessary. All items must be returned to the onsite proctor at the conclusion of the exam.

Notification of Results. Candidates are notified via email once official examination results are available within their online Candidates.ADEXtesting.org candidate portal. Generally, within 2-3 business days of exam completion.

Computer-based Exam Retakes. Eligible candidates may apply for a fee-based retake of the Local Anesthesia Computer-based Examination via the online Candidates.ADEXtesting.org candidate portal.

Official Results Reporting for Licensure Application. American Board of Dental Examiners is a testing agency, not a licensing authority. Successful completion of an examination does not constitute licensure. It is illegal to render patient treatment until a license or certificate is issued by the state or jurisdiction. Contact the respective licensing authority directly for their reporting requirements. See Candidate Guide and Scores page of ADEXtesting.org website for additional information.

Computer-based Exam References

Bassett, K.B., DiMarco, A.C., Naughton, D.K. (2015) *Local Anesthesia for Dental Professionals* (2nd ed.). Upper Saddle River: Pearson Education.

Logothetis, D.D. (2017). Local Anesthesia for the Dental Hygienist (2nd ed.) St. Louis: Elsevier.

Malamed, S.F. (2015). *Medical Emergencies in the Dental Office (7th ed.).* St. Louis: Elsevier Mosby.

Malamed, S.F. (2013). Handbook of Local Anesthesia (6th ed.). St. Louis: Elsevier Mosby.

To assist Candidates in preparing for the Local Anesthesia Computer-based (computer-based) examination, the following chart was referenced when constructing MRDs and local anesthesia dosage items.

Adult MRDs and Local Anesthesia Drug Information

Local Anesthetic Drugs	Vasoconstrictor	MRD (mg/lb)	MRD (mg/kg)	MRD ** Absolute (mg)	Duration Category	Pulpal Anesthesia (approximate minutes)	Soft Tissue Anesthesia (approximate minutes)
4% articaine * (w/ vaso)	epinephrine 1:100,000	3.2	7.0	weight dependent	Intermediate	60-75	180-300
4% articaine * (w/ vaso)	epinephrine 1:200,000	3.2	7.0	weight dependent	Intermediate	45-60	120-300
0.5% bupivacaine *	epinephrine 1:200,000	0.9	2.0	90	Long	90-180	240-540
2% lidocaine (w/vaso)	epinephrine 1:50,000	3.2	7.0	500	Intermediate	60	180-300
2% lidocaine (w/vaso)	epinephrine 1:100,000	3.2	7.0	500	Intermediate	60	180-300
3% mepivacaine (plain)	None	3.0	6.6	400	Short	20-40	120-180
2% mepivacaine (w/ vaso)	levonordefrin 1:20,000	3.0	6.6	400	Intermediate	60	180-300
4% prilocaine (plain)	None	4.0	8.8	600	Short (I) Intermediate (B)	10-15 (I) 40-60 (B)	90-120 (I) 120 - 240 (B)
4% prilocaine (w/ vaso)	epinephrine 1:200,000	4.0	8.8	600	Intermediate	60-90	180-480

Infiltration/supraperiosteal = (I)

Block = (E

Source: Adapted from Bassett, K.B. DiMarco A.C., Naughton D.K (2015) Local Anesthesia for Dental Professionals (2nd ed.) Upper Saddle River: Pearson Education

AAPD Pediatric MRDs and Local Anesthesia Drug Information

Local Anesthetic Drugs Color Coded Box	Vasoconstrictor	MRD (mg/lb)	MRD (mg/kg)	MRD Absolute (mg)	Duration Category	Pulpal Anesthesia (approximate minutes)	Soft Tissue Anesthesia (approximate
4% articaine (w/ vaso)	epinephrine 1:100,000	3.2	7.0	500	Intermediate	60	180-300
4% articaine (w/ vaso)	epinephrine 1:200,000	3.2	7.0	500	Intermediate	45	120-300
0.5% bupivacaine	epinephrine 1:200,000	0.6	1.3	90	Long	40	240-540
2% lidocaine (w/vaso)	epinephrine 1:50,000	2.0	4.4	300	Intermediate	60	190
2% lidocaine (w/vaso)	epinephrine 1:100,000	2.0	4.4	300	Intermediate	60	180-300
3% mepivacaine (plain)	None	2.0	4.4	300	Short	25 (I) 50 (B)	120-180
2% mepivicaine (w/ vaso)	levonordefrin 1:20,000	2.0	4.4	300	Intermediate	50	180-300
4% prilocaine (plain)	None	2.7	6.0	400	Short (I) Intermediate (B)	20 (I) 40-60 (B)	90-120 (I) 120 - 240 (B)
4% prilocaine (w/ vaso)	epinephrine 1:200,000	2.7	6.0	400	Intermediate	60-90	180-480

Infiltration/supraperiosteal = (I)

Block = (B)

This chart is adapted from AAPD Guidelines: Use of Local Anesthetic (2015)

^{*} No FDA weight-based recommendation, MRD= 90 mg. Health Canada weight-based recommendation 0.9 mg/lb or 2.0 mg/kg.

^{**} Dosages reflect specific local anesthetic drugs, individual absolute MRDs may also be adjusted for vasoconstrictor limits.

PATIENT-BASED CLINICAL EXAM

The purpose of the Local Anesthesia Clinical Examination is to evaluate a candidate's ability to utilize professional judgment and knowledge to competently administer both an Inferior Alveolar (IA) and a Posterior Superior Alveolar (PSA) nerve block injection on a Patient.

REMINDER. In those jurisdictions that require completion of a clinical exam for local anesthesia licensure or certification, the Local Anesthesia Examination is comprised of two components: a Computer-based exam and a Patient-based Clinical exam. All components of the Local Anesthesia Examination series must be successfully completed (passing scores) within 18 months of the initial attempt of any component of the series.

Clinical Scheduling Requests. Scheduling requests may be submitted via the "My Registration" button of your <u>Candidates.ADEXtesting.org</u> profile or via <u>Contact Us</u> prior to the **Registration Deadline** for a given exam for consideration. Scheduling requests are **not guaranteed.**

Scheduling request options may include:

- Specific clinical examination day/date
- Using a **fellow candidate (name)** as a patient. Candidates using a fellow candidate as a patient are scheduled an hour or more to accommodate both injections—yours and theirs.

Clinical Examination Schedules. Approximately one (1) month prior to the clinical exam, individual examination schedule will be posted to the <u>Candidates.ADEXtesting.org</u> online portal. Once released, exam schedule changes are not permitted.

Equipment and Materials

Armamentarium. You must provide the following:

- Aspirating syringe(s) only
- Hemostat or locking pliers.
- Blood pressure measuring device.
- Candidate and Patient appropriate protective eyewear with side shields (over-glasses or side shields, if wearing personal eyewear).

Site-Provided Equipment and Supplies. The exam site host (school) provides expendable materials for candidate use during the examination. The school is responsible for any deviation from this list. Materials provided are the brands used by the school. If you wish to use a specific brand, you must bring it with you.

Expendable Local Anesthesia Materials provided by Exam Site Host (School)					
2x2 gauze squares	Barrier covers/tape	Patient napkins/bibs			
Air/water syringe tips	Cotton-tip applicators	Sanitizing materials			
Antimicrobial mouthwash	Drinking cups	Soap			
Anesthetic(s)	Face Masks	Standard saliva ejectors			
 Local anesthetic(s) with and 	Facial Tissue	Surface disinfectant			
without vasoconstrictor	Gloves, nonlatex (S,M,L,XL)	Syringe Needles (long/short)			
 Topical anesthetic(s) 	Headrest covers	Tray covers			
Autoclave bags, small	Instrument trays, disposable	Trash bags			
Autoclave bags, medium	Paper towels				
Autoclave tape	Patient bib clips, disposable				

Do not assume that exam sites (schools) have extra instruments or equipment available for rent. Site-specific information can be found in the **Site Info Sheet** for the respective exam site host at <u>ADEXtesting.org</u>. Contact the exam site host directly with any site-specific questions regarding equipment, rental availability, etc.

Equipment Malfunctions. School personnel are available throughout the examination to resolve malfunctions of operatories and school equipment. Candidates should familiarize themselves with available school equipment during the clinic tour. Neither the American Board of Dental Examiners nor school personnel will be responsible for time lost due to the malfunction of any personal equipment.

Notify the Chief Examiner immediately of any malfunction of school equipment. An appeal cannot be based on loss of clinic time if the incident is not reported to the onsite Chief Examiner.

Patient Criteria

It is your responsibility to submit a Patient that meets the examination criteria.

All medical clearances must be written on the health care provider's letterhead, indicate clearance for the examination date range, and reflect any necessary alterations in treatment.

A Patient must meet all of the following criteria:

- 1. 18 years of age or older.
- 2. **Cannot** be a dentist, dental hygienist, or dental hygiene educator.
- 3. a) **Mandibular dentition.** At least one mandibular premolar present on the contralateral side of the submitted quadrant.
 - b) Maxillary dentition. Maxillary second molar present in the submitted quadrant.
- 4. Medical clearance if a Patient is pregnant.
- 5. Medical clearance and/or antibiotic coverage if the Patient has had a heart attack, stroke, or cardiac surgery within the past six (6) months.
- 6. Preoperative blood pressure and pulse must be recorded on the *Medical History/Patient Consent* form and taken on the Patient no more than one hour prior to the scheduled clinical examination time.
 - Systolic blood pressure of 140 or below **or** a diastolic blood pressure reading of 90 or below are acceptable. **A Patient with a systolic blood pressure reading of 141-179 **or** a diastolic blood pressure reading of 91-119 will only be accepted with written medical clearance. A Patient with a systolic reading greater than 180 and/or a diastolic reading greater than 120 is not allowed.
- 7. No orofacial herpes at the vesicle, ulcerated vesicle, or prodrome stage.
- 8. No intraoral sore or puncture mark in the two (2) potential penetration sites.
- 9. **ASA I or II.** An ASA III status that does not alter care or pose a risk to the Candidate or Patient in a testing environment.

The Patient must not have:

- 1. Active tuberculosis.
- 2. Previous Intravenous bisphosphonates therapy.
- 3. Used cocaine or methamphetamine drugs within the past 24 hours.

- 4. Any condition, medication or drug history that might be adversely aggravated by the length or nature of the examination procedures.
- 5. An Absolute contraindication to anesthetic/vasoconstrictor

Failure of the examination will result if the patient is rejected for any of the above reasons.

If your patient has a puncture mark or an intraoral sore or at the injection site, you must submit the other side.

If your initial patient has active oral herpes or high blood pressure and you are unable to submit a second patient, you will be recorded as a "No Show" Candidate. Subsequent submissions are subject to all Patient Criteria requirements for acceptance. Refer to the "No Show" Candidate section for additional information.

Blood Pressure Guidelines

Preoperative blood pressure/pulse taken and recorded on *Patient Medical History* form no more than 1-hour prior to scheduled clinical examination.

CONDITION	BLOOD PRES	SURE (and/or)	ACTION	
CONDITION	Systolic	Diastolic	ACTION	
Normal	< 120	< 80		
Elevated	120-129	< 80	Acceptable to continue with treatment	
Stage 1 Hypertensive	130-139	80-89		
Stage 2 Hypertensive	<u>≥</u> 140	<u>≥</u> 90	Acceptable to Continue ONLY with written consent from Patient's health care provider	
HYPERTENSIVE CRISIS (Emergency care needed)	>180	>120	Unacceptable for treatment	

Source: US Department of Health and Human Services. National Institutes of Health. National Heart, Lung and Blood Institute. National High Blood Pressure Education Program. www.nhlbi.nih.gov

Malpractice Insurance. Candidate registration for the patient-based clinical exam includes limited professional liability coverage during the examination. Additional limited liability insurance is not required.

Grading Criteria

You will be expected to perform both injections within 20 minutes. If you are unprepared, or taking a considerable amount of time, the examiners will remind you of the time restraints.

Critical Aspects and Less Critical Aspects of Injection

Each aspect of the injection is classified as **critical** or **less critical**. Eight (8) critical aspects of the injection have an asterisk (*). One (1) critical aspect error validated by both examiners, results in failure of the examination. Three (3) less critical aspect errors in Category 2 (validated by both Examiners) results in failure of the examination.

You may change your submission, without penalty, at any time prior to an examiner checking landmark teeth and penetration sites.

*1. Medical History, Oral Conditions, Anesthetic and Syringe Selection

Medical History

Errors:

· See Patient Criteria

Oral Conditions

IA Injection Errors:

- Missing both mandibular premolars on the contralateral side
- · Intraoral sore or puncture mark at the injection site
- Orofacial herpes is failure of both injections

PSA Injection Errors:

- Missing maxillary second molar in the submitted quadrant.
- · Intraoral sore or puncture mark at the injection site
- Orofacial herpes is failure of both injections

Anesthetic & Syringe Selection

Errors:

- Long-acting anesthetic
- · High concentration vasoconstrictor
- Self-aspirating or non-threading syringe

2. Syringe Preparation and Handling

Armamentarium

Errors:

- Appropriate protective eyewear is not worn by the candidate and/or patient
- Locking pliers or hemostat not present
- Expired local anesthetic
- Anesthetic selection or needle length does not match Local Anesthesia Information Sheet

Syringe Properly Prepared

Errors:

- Harpoon is not securely engaged
- Large bubbles not expelled from cartridge prior to injection
- Expelled solution is more than stopper width
- · Needle not securely attached to the syringe

Syringe Handling

Error:

Syringe in patient's field of vision prior to injection

*3. Penetration Site

Needle Contamination

Error:

• Needle touches any surface, facial anatomy or intraoral object (gauze, glove) prior to penetration. Proceeding with a contaminated needle results in failure of the injection.

Three Penetrations Allowed

Penetration of the tissue is considered an attempt. Three (3) penetrations are allowed to obtain the optimum angle and depth. If the third penetration results in a first positive aspiration, a fourth penetration is permitted.

Error:

Optimum depth and angle is not obtained in three (3) penetrations

Optimum IA Nerve Block Penetration Site

The penetration site is at the area bordered medially by the pterygomandibular raphe, laterally by the internal oblique ridge, and at the height of the coronoid notch.

Errors:

- Too superior
- Too inferior
- Too medial
- Too lateral

Optimum PSA Nerve Block Penetration Site

The penetration site is at the height of the vestibule in the mucobuccal fold, posterior to the zygomatic process of the maxilla (visually, this approximates the distal facial root of the 2nd molar).

Errors:

- Too anterior
- Too posterior
- Not in the mucobuccal fold

*4. Angle and Depth

Optimum IA Angle and Depth

At the optimum depth, the barrel of the syringe should be positioned over the premolars on the contralateral side and the needle parallel to the occlusal plane of the mandibular teeth.

The depth of insertion is 20 mm - 25 mm (approximately 2/3 the length of long needle or 4/5 the length of short needle).

Errors:

- Barrel too distal
- Barrel too mesial
- · Angle too high
- Angle too low
- Too shallow
- Too deep

Optimum PSA Angle and Depth

At the optimum depth, the barrel of the syringe must be positioned 45 degrees to the occlusal plane and 45 degrees to the midline.

The depth of insertion is approximately 16 mm (about 1/2 the length of a long needle or 3/4 the length of a short needle).

Errors:

- Barrel of the syringe not at 45-degree angle toward midline and/or barrel of the syringe not at 45-degree angle to occlusal plane
- Too shallow
- Too deep

*5. Aspiration

Large Window Visible

Upon aspirating, the large window must be toward the Candidate.

Error:

· Small window toward the candidate

Aspiration Observed

Errors:

- No visible movement of the stopper upon aspiration
- · Harpoon is not engaged after aspiration is declared to be negative

Proper Handling of Positive Aspiration

A positive aspiration is observed and handled correctly.

Errors:

- Proceeding after aspiration when a significant amount of blood is present in the cartridge
- Failing to recognize a positive aspiration

*6. Amount and Rate

Deposition of Anesthetic Prior to Aspiration

Frror

• More than 1/4 of the anesthetic is deposited in the tissue, prior to reaching the optimum depth and before aspiration.

Rate of Administration is Acceptable

Acceptable rate of depositing anesthetic is approximately fifteen (15) seconds for two (2) stopper widths.

Error:

 Rate of deposition is approximately 12 seconds or less for two (2) stopper widths of anesthetic.

*7. Tissue Management

Tissue Management is observed throughout the administration of the injection.

Errors:

- Needle is visibly bent upon removal from tissue
- Degree of the bowing would likely result in excessive submucosal soft tissue injury
- Visible laceration of tissue upon approach or withdrawal of the needle with or without bleeding
- Safety and well-being of the patient is compromised

*8. Recapping

Recapping is observed throughout the examination.

Proper Recapping Technique

A one-handed recapping technique is required. Once the needle is safely inside the needle cap (protected), the cap must be secured.

Errors:

- A two-handed recapping technique is observed during recapping
- Holding the needle cap with a hemostat or locking pliers

*9. Sharps Disposal

All needles and cartridges must be disposed of properly and according to exam site host (school) policy. Sharps Disposal is evaluated after the completion of both injections.

Errors:

- Sharps and cartridges are not properly disposed of according to exam site host (school) policy
- · Two-handed recapping is observed

Improper Sharps Disposal will result in failure of both injections.

Stop and Hold/Failure of the Injection

When any procedure poses a health or safety risk to the Patient a Stop and Hold is announced. This signifies that a **critical error** has occurred and results in failure of that injection.

You must stop and hold your position. If the needle has penetrated the tissue, you will be instructed to withdraw.

The following are reasons for Stop and Hold:

- Absolute contraindication to anesthetic/vasoconstrictor
- Long-acting anesthetic
- High concentration vasoconstrictor
- Needle Contamination
- Three Unsuccessful Penetrations
- Small window toward candidate upon aspiration
- No visible movement of the stopper
- Harpoon is not engaged upon aspiration
- Improper handling of a positive aspiration
- More than ¼ of the anesthetic is deposited prior to reaching deposition site (for depth and angle)
- Rate of deposition is too rapid
- Improper Tissue Management
- Improper Recapping
- Improper Sharps Disposal

Clinical Examination Day

Anonymity of the Examination. The Local Anesthesia clinical examination is administered by examiners and is not an anonymous examination. However, your name or school identification must not appear on any of the examination materials, equipment, instruments or clothing.

Onsite Question and Answer Session/Clinic Tour

An onsite Question and Answer (Q&A) Session is held prior to your scheduled examination. The Chief Examiner or Exam Technical Coordinator (ETC) will verify candidate identification and distribute exam materials.

The following forms are provided during the onsite Q&A Session:

- Patient Medical History/Patient Consent form
- Local Anesthesia Injection Information Sheet

The Clinic Tour is optional and will familiarize you with the exam site host (school) clinic layout, emergency protocols, infection control policies, proper disposal of biohazardous and pharmaceutical materials, and operation of the equipment.

To expedite the flow of the examination, complete the Patient Medical History/Patient Consent form prior to clinic entrance.

"No Show" Candidate. You may opt to withdraw from the exam and forfeit your examination attempt and registration fees for the following reasons:

- You do not have required items or are not dressed in proper clinic attire at your assigned clinic time.
- · You are late for your assigned clinical examination.
- Your first patient has been rejected for active oral facial herpes or high blood pressure.

Withdrawal for any other reason constitutes failure of the examination.

Clinic Entrance and Exam Procedures

Arrive onsite and be prepared 30 minutes prior to your scheduled time. Candidates will be notified when to enter the clinic and directed to their operatory. After setting up the operatory, bring your patient back.

If you did not attend the Q&A Session, the Chief Examiner will verify your identification and distribute your exam materials prior to clinic entrance.

An Examiner will check the following:

- Patient Medical History/Patient Consent form
- Two penetration sites
- · Landmark teeth are present in each submitted quadrant
- Locking pliers or hemostat present
- Syringe selection correct
- · Patient and Candidate eyewear present
- · Anesthetic selection matches Local Anesthesia Information Sheet

No more than two prepared syringes may be present on your tray. (Additional cartridges and needles can be set aside on a non-contaminated surface to load.) The anesthetic type and expiration date on the cartridge should be visible for Examiner review. Do not loosen the needle cap(s) until directed by the Chief Examiner.

Examiners will instruct you when to apply topical; however, topical anesthetic use is at the discretion of the patient.

The inferior alveolar (IA) nerve block injection will be administered first. The inferior alveolar (IA) nerve block and posterior superior alveolar (PSA) nerve block may be performed on the same side or opposite side of the mouth.

Two (2) examiners will observe your technique. Both injections must be performed to criteria to pass the clinical examination. (You may describe your actions as the injection is being performed.)

Pause and Announce. There are four (4) times that you must pause and announce each critical phase of the injection.

- 1. **Initial Penetration.** After the needle has penetrated the tissue, stop and hold the position. Wait until instructed to proceed.
- 2. **Angle and Depth.** Advance to the deposition site. Inform the Examiners when you have reached the optimum depth and angle. Wait until instructed to proceed.
- 3. **Aspiration.** Announce when aspirating and whether the aspiration is positive or negative. If the aspiration is negative, wait until instructed to proceed.

If the aspiration is positive, an Examiner will ask," How you would like to proceed?" Use your professional judgment in determining how to handle the situation. A positive aspiration on two (2) attempts does not automatically result in failure.

NOTE: Aspiration is required on only one (1) plane. There is no penalty if an aspiration is completed on two (2) planes.

4. **Deposition Rate.** Inform the examiners prior to depositing anesthetic. You will be instructed when to withdraw.

Your technique is evaluated at each critical aspect. After observing each critical aspect, one examiner will say "I see" and the other examiner will state, "Proceed."

Upon completion of the IA nerve block, secure the cap or proceed to the PSA nerve block.

Exam Completion

After completion of both injections, perform the following:

- 1. Properly dispose of contaminated sharps and cartridges. Refer to exam site host (school) policy regarding unused anesthetic cartridges.
- 2. Record on the *Anesthesia Information Sheet*, the total volume of local anesthetic (in milliliters or milligrams) administered.
- 3. Dismiss your patient, remove barriers, disinfect operatory and exit the clinic.
- 4. Return any unused Candidate ID Labels and any applicable *ITCs* to the designated location.

Preliminary Results Reporting

Candidates are notified of their preliminary examination **results via text** (844). If not received, check the "spam" filter of your SMS/RCS text messaging app. Or edit the phone number displayed in the "My Account–Personal Info" section of your online <u>Candidates.ADEXtesting.org</u> profile. If necessary, contact the onsite ETC to resend the preliminary results text.

If unsuccessful, any applicable criterion (reason for failure) will be displayed. Discussion with examiners or staff regarding performance or reasons for failure is prohibited.

To register for a same-day onsite retake exam, utilize the "**Schedule Retake - Today**" link provided in your preliminary examination results text. Complete payment using a valid credit or debit card—Visa or Mastercard. Once complete, contact the onsite ETC to receive your assigned retake time and examination forms.

Optional Onsite Retake Exam

An onsite fee-based retake is available during the examination for eligible candidates. A retake is considered a separate examination and included in your total number of examination attempts.

Candidates are permitted up to four (4) unsuccessful attempts of the Clinical Examination at two (2) separate exam locations prior to remediation (ex. Clinical + Retake = two attempts).

If you elect to retake the exam at the same location, you will perform only the failed injection(s). If you re-attempt the examination at another location, you will be required to perform both IA and PSA nerve block injections, and pay both the registration fee and facility fee.

Performance Evaluation. Exam criteria and scoring are the same.

Patient Criteria Options. Your same patient or a new patient may be submitted. When submitting a previous patient, the injection(s) must be performed on the opposite side.

If an unsuccessful examination attempt occurred prior to penetrating the tissue (i.e., needle contamination) the retake may be performed on the initial side.

Notification of Results. Candidates are notified via email once official examination results are available within their online Candidates.ADEXtesting.org candidate portal. Generally, within 2-3 business days of exam completion.

Official Results Reporting for Licensure Application. The American Board of Dental Examiners is a testing agency, not a licensing authority. Successful completion of an examination does not constitute licensure. It is illegal to render patient treatment until a license or certificate is issued by the state or jurisdiction. Contact the respective licensing authority directly for their reporting requirements. See <u>Candidate Guide</u> and <u>Scores</u> page of <u>ADEXtesting.org</u> website for additional information.

SAMPLE FORM

	PAT	IENT CONSENT AND AS	SUMPTION OF R	ISK			
	LOCAL ANESTHESIA PATIENT MEDICAL HISTORY CANDIDATE IS PATIENT FIRST NAME						
DATE OF BIRTI Circle "YES" of Do you have or A B C D A B C D A A A A A A A A A A A A		I Tuberculosis J Kidney/Renal Dis K Hepatitis/Jaundio L HIV Positive M Epilepsy/Seizure N Joint Replaceme D Liver/Hepatic Dis P Latex Allergy	yes No YE		ITA and the reating you ing a dental amination to ITA state. you receive examiners) / up dental abstandard, inistered by I expressly due to the The School rided by the		
4. Are you cu5. Within the physician o6. Have you c7. Have you u	rently receiving or have you previously ast six months, have you been seen but health care provider? experienced local anesthetic complicates any recreational drug(s) (cocaine the or have your been exposed to any complete the or have your been exposed	y received intravenous bisphos y, or are you currently under th ons with dental treatment in th or methamphetamines) within	e care of a e past? the last 24 hours?	YES NO	or after the rs) and the examiners) demands, , nature or ng or after on of Risk. I anesthetics and that my exam and		
LOCAL A Patient First Name: Initial Exam Attempt	Onsite Retake Exam Atten	npt		lical clearance	2024		
	Date:	FIFNOTH					
INJECTION II		SHORT					
PSA LEI	37-0. 90-00-00-00-00-00-00-00-00-00-00-00-00-0	SHORT					
Complete this section after eva	luation; form remains at operatory.	0.0000000000000000000000000000000000000					

Visit <u>ADEXtesting.org</u> to download full-size examination forms.

Patient-Based Clinical Examination Overview

Be onsite (approximately 30 minutes prior to your assigned clinic time)

Clinic Entrance

- Patient Medical History/Patient Consent form completed (vitals taken one-hour or less prior to exam)
- · Local Anesthesia Information Sheet completed
- Set up operatory
- Prepare syringe(s) with chosen local anesthetic and needles

Clinical Examination Preparation

One examiner will check:

- Penetration sites are acceptable and landmark teeth are present
- Syringes are properly prepared
- Local Anesthesia Information Sheet is accurate
- · Medical History and patient vitals reviewed

Examiners will inform you when to:

- Apply topical
- Loosen needle cap(s)
- Proceed with Sharps Disposal
- Dismiss your patient

After Sharps Disposal

- Record total volume (ml or mg) of local anesthetic on *Local Anesthesia Information*Sheet (form remains at operatory)
- Clean and disinfect operatory
- Return any unused Candidate ID Label to designated area