PROSTHODONTIC CROWN PREPARATION PREMOLAR ZIRCONIA

ADEX 2026

Criteria applicable August through December 31, 2026

CRITICAL ERRORS		
Wrong tooth/surface treated	NO	YES
Procedure not challenged	NO	YES

NOTE: Those SUBs that are highlighted are part of the 3-SUB Rule

ACC = Adheres to Criteria	SUB = Marginally Substandard	DEF = Critical Deficiency
---------------------------	------------------------------	----------------------------------

CERVI	CERVICAL MARGIN AND DRAW				
Margin/Extension					
ACC	The cervical margin is \leq 0.5 mm below [apical to] the simulated free gingival margin to \leq 1.5 mm above [occlusal to] the simulated free gingival margin.				
SUB	The cervical margin is over-extended > 0.5 mm below the simulated free gingival margin.				
DEF	 A. The cervical margin is over-extended by > 0.5 mm below the simulated free gingival margin, causing visual damage to the typodont. B. The cervical margin is under-extended by > 1.5 mm above the simulated free gingival margin. 				
Margi	Margin/Definition/Unbeveled				
ACC	The cervical margin is continuous but may be slightly rough and may lack some definition.				
DEF	A. The cervical margin has no continuity or definition. B. The cervical margin is beveled. C. The cervical margin is cupped or J-shaped.				
Margin Width					
ACC	The cervical margin is > 0.5 mm but ≤ 1.5 mm in width.				
SUB	The cervical margin is > 1.5 mm but ≤ 2.0 mm in width.				
DEF	A. The cervical margin is > 2.0 mm in width. B. The cervical margin is < 0.5 mm in width.				
Line o	f Draw				
ACC	The path of insertion/line of draw deviates < 20° from the long axis of the tooth.				
SUB	The path of insertion/line of draw deviates 20° to < 30° from the long axis of the tooth.				
DEF	The path of insertion/line of draw deviates ≥ 30° from the long axis of the tooth.				
WALLS	S, TAPER AND LINE ANGLES				
Axial 1	Tissue Removal				
ACC	The axial tissue removal is ≥ 1.0 mm but ≤ 2.0 mm.				
SUB	The axial tissue removal is > 2.0 mm but ≤ 2.5 mm.				
DEF	A. The axial tissue removal is > 2.5 mm. B. The axial tissue removal is < 1.0 mm.				
Axial \	Axial Walls Smoothness/Undercut				
ACC	The walls may be slightly rough and may lack some definition.				
DEF	There is an undercut, which, when blocked out, would compromise margin width criteria and/or is > 0.5 mm deep.				

Taper			
	T : 15 1 11 11 14 14 20 11		
ACC	Taper is present, from nearly parallel to ≤ 12° per wall.		
SUB	There is excessive taper that is > 12° and ≤ 16° per wall.		
DEF	Taper is grossly over-reduced > 16° per wall.		
Occlus	Occlusal Reduction		
ACC	Occlusal reduction is ≥ 1.0 mm but ≤ 2.0 mm.		
SUB	Occlusal reduction is > 2.0 mm but ≤ 2.5 mm.		
DEF	A. Occlusal reduction is > 2.5 mm. B. Occlusal reduction is < 1.0 mm.		
Internal Line Angles			
ACC	Internal line angles and cusp tip areas may not be completely rounded and may show a slight tendency of being sharp.		
DEF	The internal line angles or cusp tip areas are excessively sharp with no evidence of rounding.		
TREATMENT MANAGEMENT			
Condi	Condition of Adjacent/Opposing Teeth		
ACC	Any damage to adjacent tooth/teeth can be removed with polishing without adversely altering the shape of the contour and/or contact.		
SUB	A. Damage to adjacent tooth/teeth requires recontouring that changes the shape and/or position of the contact. B. Opposing hard tissue shows minimal evidence of damage and/or alteration inconsistent with the procedure.		
DEF	A. There is gross damage to adjacent tooth/teeth requiring a restoration. B. There is evidence of gross damage and/or alteration to opposing hard tissue inconsistent with the procedure.		
Condi	Condition of Surrounding Tissue		
ACC	There may be slight damage to simulated gingiva and/or typodont consistent with the procedure.		
SUB	There is iatrogenic damage to the simulated gingiva and/or typodont inconsistent with the procedure.		
DEF	There is gross iatrogenic damage to the simulated gingiva and/or typodont inconsistent with the procedure.		